CUMULATIVE AUTHOR INDEX
Annals of the Entomological Society of America
Volume 103, 2010

A

Abe, Y. 322, 838
Alexandrakis, V., 66
Alvarez, N., 742
Al-Wahaibi, A. K., 236
Andersen, J. C., 30, 844
Andrade-Souza, V., 424
Anfora, G., 813
Aoki, K., 379
Arbogast, R. T., 389
Audenaert, B., 247
Axen, H. J., 11
Ayabe, Y., 806

B

Bagis, N., 943
Bardunias, P., 449
Barry, K. L., 120
Başibüyük, H. H., 625
Bento, J.M.S., 742
Bergh, J. C., 50, 993
Bernard, E. C., 472
Bhatt, S., 227
Birouraki, A., 943
Blythe, L. D., 11
Boettner, G. H., 135
Bonte, M., 597
Bourtzis, K., 1003
Brent, C. S., 300
Briano, J. A., 71, 635
Britten, H. B., 1012
Brlansky, R., 827
Brooks, T., 592
Brunel, O., 111, 430
Buckley, Y. M., 196

C

Cabrera, S. M., 71, 635
Calcaterra, L. A., 71, 635
Callahan, J., 135
Campbell, L. A., 925
Campbell, N. K., 135
Carpenter, J. E., 611
Castaño-Meneses, G., 7
Castle, del conte S., 592
Caterino, M. S., 785
Chanchao, C., 965
Chang, H., 866
Chatzimanolis, S., 785
Chen, X.-X., 360
Chmielewski, M. W., 734
Cicero, J. M., 979
Cid, M., 404
Condon, M. A., 11
Contreras-Ramos, A., 695
Cook, L., 196
Correia, E. T., 603
Costa, M. A., 424
Cuezzo, F., 71, 635
Cutwa, M., 757

D

Danka, R. G., 146
De Barro, P., 196
De Bruyn, L., 247
De Clercq, P., 597
de León, J. H., 100
de Remes Lenicov, A. M., 332
Delabie, J.H.C., 424
Delaplane, K. S., 711
Dinsdale, A., 196
Dodds, K. J., 39
Donahue, C. P., 135
Dou, W., 661
Dougherty, K. M., 683
Drosopoulou, E., 1003
Duso, C., 165

E

Ebert, T. A., 827
Edde, P. A., 796
Elkinton, J. S., 135
Ellis, J. D., 711
Emlen, D. J., 209
Emoto, J., 988
Er, A., 273
Ergin, E., 273
Escher, R. L., 757
Evans, E. W., 750
Evans, J. D., 671
Evans, W. G., 823

F

Fathipour, Y., 574
Feder, J. L., 908
Felderhoff, K. L., 472
Feng, H., 875
Fereres, A., 404
Fernandes, F. S., 603
Fisher, J. R., 548
Fombong, A. T., 389
Forbes, A. A., 908
Frank, D. L., 50, 993
French, B. W., 688
French, J. V., 898
Fujisaki, K., 439

G

Galante, E., 956
Gálvez, S., 96
Gammons, J. R., 11
Gao, Y., 7
Garcia-Medina, C., 544
García-López, A., 956
Gassmann, A. J., 371
Gilmore, D. W., 39
Gingras, S., 592
Goolsby, J. A., 252
Greene, K., 757
Gruwell, M. E., 844
Gwiazdowski, R., 135

H

Haight, K. L., 678
Hall, D. G., 611
Hamm, C. A., 20
Hammack, L., 688
Handler, A. M., 264
Hansen, L. D., 418
Harris, J. W., 146
Harrison, J. L., 11
Havelka, J., 307
He, J.-H., 360
Hernández, M. C., 332
Herrmann, D. L., 227
Hoelmer, K. A., 532
Hoffmann, A. A., 886
Hong, X.-Y., 697
Hood, G. R., 908
Hood, W. M., 671, 711
Hu, F., 661
Hunkins, R. R., 135

I

Ibañez, J., 96
Ide, T., 838
Ioriatti, C., 813
Ishizaki, M., 585

J

James, D. G., 592
Jannot, J. E., 227
Jia, F.-X., 661
Johansen, K. J., 548
Johns, P. M., 120
Jones, G. D., 566
Jourdie, V., 742
Juliano, S. A., 227, 757, 1038

K

Kajita, Y., 750
Karimzadeh, J., 574
Kato, M., 379
Kavallieratos, N. G., 153, 307
Kawakami, Y., 327

Keena, M. A., 949
Kesavaraju, B., 1038
Khatchikian, C., 734
Khrimian, A., 135
Kim, H., 532
Kirejtshuk, A., 866
Kishi, S., 580
Klaric, V., 165
Ko, A. E., 227
Koeppler, K., 1003
Komazaki, S., 916
Kontodimas, D. C., 943
Korkmaz, E. M., 625
Kounatidis, I., 1003
Kreiter, S., 165
Kwon, Y.-D., 532

L

Landolt, P., 592
Lauzière, I., 57
Law, Y.-H., 771
Lee, C.-F., 500
Lee, S., 532
Lee, W., 532
Lee, Y.-F., 216
Leimkuehler, W., 84
Leopold, R. A., 264
Leskey, T. C., 50, 993
Lewis, S. M., 128
Li, C.-L., 341
Li, Y.-P., 647
Lin, Y.-H., 216
Liu, C., 875
Liu, C.-Y., 1018
Liu, Y.-Q., 647
Livdahl, T., 734
Lounibos, L. P., 757
Lounsberry, Z., 671
Louzada, E., 898
Lu, C., 647
Lucchi, A., 813
Luna, M. G., 936
Lynch, C., 1012

M

Makarkin, V. N., 511, 855
Malaquias, J. B., 603
Mariano, C.S.F., 424
Mata, J., 898
Matsumoto, T., 716
Mattos, I., 523
Mavragani-Tsipidou, P., 1003
Maxwell, M. R., 120
Mazzoni, V., 813
McNish, I. G., 11
Medianero, E., 492
Meinke, L. J., 925
Mermudes, J.R.M., 523
Meza, J. S., 96
Mó, E., 956
Miller, C. W., 209
Miller, R. S., 775
Ming, Q.-L., 128
Miura, K., 366
Mlynarek, J. J., 465
Moe, A. M., 1025
Mogi, M., 618
Monroe, E. M., 1012
Moran, P. J., 252
Morse, G. E., 844
Morse, J. G., 30, 236
Moulton, J. K., 472
Moya-Raygoza, G., 544
Muli, E., 389
Murakami, N., 379
Murillo, H., 742
Mutyambai, D. M., 389

N

Nagoshi, R. N., 283
Nakajima, Y., 439
Nakou, I., 1003
Nascimento Júnior, J. L., 603
Neumann, P., 671
Neven, L. G., 418
Nicácio, J., 723
Nieves-Aldrey, J. L., 492
Nishida, T., 580, 716
Nishimura, N., 757
Noriyuki, S., 580, 716
Normark, B. B., 30, 844
Norris, L. A., 785
Numa, C., 956

O

Oku, K., 802
Okutani-Akamatsu, Y., 802
O'Meara, G. F., 757
O'Neill, K. M., 775
O'Neill, R. P., 775
Ornelas, J. F., 397

P

Palacios-Vargas, J. G., 7
Papadopoulos, N. T., 1003
Pearce, A. M., 775
Peck, S. B., 1
Peréz, I. J., 635
Pernal, S. F., 671
Petroć-Obradoć, O., 153
Pettis, J., 671
Phillips, P. K., 860
Phillips, T. W., 796
Pitts, J. P., 555
Plague, G. R., 683
Porter, A. H., 135
Pucci, T. M., 566
Pultar, O., 949

Q

Qin, L., 647

R

Rajamohan, A., 264
Rakhshani, E., 307
Ramalho, F. S., 603
Rattanawannee, A., 965
Reiskind, M. H., 757
Ren, B., 654
Ren, D., 855, 866, 875
Reyna, V., 592
Ridland, P. M., 886
Riginos, C., 196
Rivers, D. B., 273
Robinson, A. S., 96
Rogers, M., 827
Rugman-Jones, P. F., 30
Rull, J., 111, 430

S

Sak, O., 273
Sakuma, M., 439
Sammataro, D., 979
Sanborn, A. F., 860
Santos, I. S., 424
Sánchez, N. E., 936
Sari, M., 625
Sasaki, R., 439
Schaefer, C. W., 706
Schal, C., 84
Scheufele, S. B., 135
Schultz, T. R., 181
Sediqi, A., 771
Sétamou, M., 57, 100, 898
Seybold, S. J., 39
Shang, L., 654
Shapiro, J. P., 611, 971
Sharkey, M., 307, 548
Shatters, R. G., 611
Shelly, T. E., 264
Shi, S.-L., 647
Shigehara, T., 916
Shih, C., 875
Shirk, P. D., 971
Silva, J. G., 424
Singer, M. S., 371
Soluk, D. A., 1012
Song, Z.-W., 697
Sonstegard, T. S., 671
Sosa-Calvo, J., 181
Soufbaf, M., 574
Spiewok, S., 671
Sremac, M., 135
Starý, P., 153, 307
Stock, S. P., 371
Stouthamer, R., 30
Su, N.-Y., 449

T

Tabashnik, B. E., 371
Takabayashi, J., 366
Tan, J.-L., 360
Tatsuta, H., 327
Teal, P.E.A., 389
Tixier, M.-S., 165
Toda, S., 916
Tomalski, M., 84

Tomanović, Z., 153, 307
Torto, B., 389
Tsitsipis, I., 66
Turlings, T.C.J., 742

U
Uchôa, M. A., 723
Uçkan, F., 273

V
Valenzuela, I., 886
Van Gossum, H., 247
Vandel, A., 949
Vargo, E. L., 84
Varikou, K. N., 66, 943
Velázquez, T., 397
Villa, J. D., 146
Virant-Doberlet, M., 813
Virla, E., 742
Voltaire, G., 683

W
Wachi, N., 322, 838
Wada, V. I., 936
Walsh, B. E., 683
Wang, C.-C., 341
Wang, J.-J., 661
Wang, P., 654
Wang, S., 654
Wang, Y., 654
Watanabe, H., 293
Watanabe, T., 585, 802
Weeks, A. R., 886
Weiblen, G. D., 1025
Wheeler, T. A., 465
Whited, B. M., 135
Wilson, J. S., 555
Winterton, S. L., 511
Wongsiri, S., 965
Wright, L. C., 592
Wu, I.-H., 1018
Wu, S., 647
Wu, S.-H., 216

X
Xia, R.-X., 647
Xiang, Z.-H., 647
Xue, X.-F., 697

Y
Yamashita, K.-I., 366
Yang, P.-S., 341, 1018
Yang, Q., 855
Yano, E., 293
Yasuda, T., 585
Yeh, W.-B., 1018
Yoshida, A., 988

Z
Zack, R. S., 706
Zalucki, M. P., 574
Zanuncio, J. C., 603
Zepeda-Cisneros, C. S., 96
Zhang, G.-N., 661
Zhang, J., 455
Zhao, Y., 875
Zikić, V., 153

CUMULATIVE KEY WORD INDEX
Annals of the Entomological Society of America
Volume 103, 2010

A

Abagrotis orbis,
  Agrotis vetusta, 592
  cutworm, 592
  grape, 592
  Noctuidae, 592
Abgrallaspis,
  armored scale insects, 30
  avocado, 30
  insect pest, 30
  molecular phylogenetics, 30
Acarapis woodi,
  Apis mellifera, 711
  colony decline, 711
  varroa mite, 711
acoustic behavior,
  cicadas, 860
  morphometrics, 860
  systematics, 860
  thermal adaptation, 860
adaptation of populations,
  Hemiptera, 544
  leafhoppers, 544
  maize, 544
Aedes,
  Culicidae, 618
  life history trait, 618
  Nepenthes pitcher plant, 618
  Stegomyia, 618
Aethina tumida, 389, 671
  Apis mellifera, 671
  DNA barcode, 671
  honey bee, 389
  invasive species, 671
  Kenya, 389
  Kodamaea ohmeri, 389
  Oplostomus haroldi, 389
  pollination, 671
Africa,
  Elachipterini, 465
  Oscinellinae, 465
  phylogeny, 465
  taxonomy, 465
agave,
  mating behavior, 430
  natural history, 430
  Tephritoidea, 430
  Ulidiidae, 430
age,
  female mating receptivity, 802
  male courtship, 802
  Miridae, 802
  ovarian development, 802
age-specific mortality,
  death frequency, 574
  Plutella xylostella, 574
aggregation pheromone,
  beetle density, 796
  lesser grain borer, 796
  nutritional stress, 796
  pheromone output, 796
agricultural pest,
  mitotic chromosomes, 1003
  polytene chromosomes, 1003
  Rhagoletis completa, 1003
  Wolbachia pipientis, 1003
Agrotis vetusta,
  Abagrotis orbis, 592
  cutworm, 592
  grape, 592
  Noctuidae, 592
alkaloids,
  entomopathogenic nematodes, 371
  pathogen reproduction, 371
  polyphagy, 371
  trade-off, 371
allometry,
  genetic differentiation, 209
  microevolution, 209
  phenotypic plasticity, 209
  sexual selection, 209
alternation of generations,
  eastern Palearctic, 838
  new species, 838
  oak-gall wasp, 838
  ring-cupped oak, 838
Alydinae,
  generic description, 706
  Leptocorisini, 706
  Melanacanthus, 706
  Micrelytrinae, 706
amplified fragment length polymorphism,
  cytochrome oxidase I, 1018
  genetic diversity, 1018
  Troides aeacus, 1018
Anastrepha ludens, 96, 264
  Anastrepha suspensa, 264
  Ceratitis capitata, 264
  fitness testing, 264
  inheritance, 96
  linkage analysis, 96
  long-term cold storage, 264
  mutations, 96
Anastrepha suspensa,
  Anastrepha ludens, 264
  Ceratitis capitata, 264
  fitness testing, 264
  long-term cold storage, 264
Anastrepha spp.,
  Ceratitis capitata, 723
  host fruit, 723
  Neosilba spp., 723
  taxonomy, 723
Andricus,
  heterogony, 322
  life cycle, 322
  synonym, 322
antennal morphology, 661, 993
  olfaction, 993
  scanning electron microscopy, 661
  sensilla, 661, 993
  Synanthedon scitula, 993
  Tephritidae, 661
antennal sensilla,
  Bombus, 654
  rough set, 654
Antheraea pernyi,
  cloning, 647
  expression pattern, 647
  lysophospholipase, 647
Anthribinae,
  fungus weevils, 523
  morphology, 523
  phylogeny, 523
  Ptychoderini, 523
antipredatory behavior,
  container community, 1038
  predation risk, 1038
Aphidiinae, 153, 307
  aphids, 153
  Cucurbitaceae, 153
  Greenideinae, 307
  key, 307
  review, 307
  Solanaceae, 153
  southeastern Europe, 153
aphids,
  Aphidiinae, 153
  Cucurbitaceae, 153
  Solanaceae, 153
  southeastern Europe, 153
Aphis glycines,
  Aphis gossypii, 532
  host alternation, 532
  molecular identification, 532
  morphology, 532
Aphis gossypii,
  Aphis glycines, 532
  host alternation, 532
  molecular identification, 532

morphology, 532
Apis mellifera, 671, 711, 979
Acarapis woodi, 711
Aethina tumida, 671
colony decline, 711
crop, 979
cuticle, 979
DNA barcode, 671
honey bee, 979
honey stomach, 979
invasive species, 671
pollination, 671
varroa mite, 711
Apis,
geometric morphometry, 965
honey bee, 965
wing morphometry, 965
Appalachian Mountains,
Collembola, 472
phylogeny, 472
taxonomy, 472
Tomoceridae, 472
armored scale, 252, 844
biological control, 252
biology, 252
cryptic species, 844
Diaspididae, 844
giant reed, 252
non-native invasive grass, 252
oleander scale, 844
species delimitation, 844
armored scale insects,
Abgrallaspis, 30
avocado, 30
insect pest, 30
molecular phylogenetics, 30
arrest of reproduction,
diapause, 366
Leptocorisa chinensis, 366
recovery of reproduction, 366
artificial diet, 597, 949
biological control, 597
Lymantria monacha, 949
mating, 597
Orius laevigatus, 597
phenology, 949
rearing, 949
reproduction, 597
Ascidae,
fecundity, 397
flower mites, 397
pollen feeding, 397
Asian citrus psyllid,
biological control, 100
DNA fingerprinting, 100
ectoparasitoid, 100
natural enemy, 100
Asopinae,
biology, 603
natural enemy, 603
photophase, 603
stink bug, 603
Asteraceae,
mouthpart length, 566
nectar feeding, 566
parasitic wasps, 566
ATP,
codling moth, 418
controlled atmospheres, 418
differential scanning calorimetry, 418
metabolism, 418
avocado,
Abgrallaspis, 30
armored scale insects, 30
insect pest, 30
molecular phylogenetics, 30
Axymyiidae,
Diptera, 455
Jurassic, 455
male adults, 455
taxa, 455

B

barcoding,
cytochrome oxidase 1, 196
phylogenetics, 196
whitefly, 196
Bayesian inference,
internal transcribed spacer, 742
mitochondrial DNA, 742
parasitoids, 742
spatial genetic structure, 742
beetle density,
aggregation pheromone, 796
lesser grain borer, 796
nutritional stress, 796
pheromone output, 796
behavioral choice,
Florida, 227
local adaptation, 227
oviposition choice, 227
phenotypic variation, 227
biodiversity,
biological conservation, 71
community structure, 71
faunal survey, 71
functional groups, 71
biogeography, 360, 555
desert, 555
Minangini, 360
Oriental, 360
Palearctic, 360
phylogeography, 555
Sigalphinae, 360
Sphaeropthalminae, 555
velvet ants, 555
biological conservation,
biodiversity, 71
community structure, 71
faunal survey, 71
functional groups, 71
biological control, 100, 236, 252, 597
armored scale, 252
artificial diet, 597
Asian citrus psyllid, 100
biology, 252
DNA fingerprinting, 100
ectoparasitoid, 100
giant reed, 252
glassy-winged sharpshooter, 236
Homalodisca liturata, 236
mating, 597
morphometrics, 236
natural enemy, 100
non-native invasive grass, 252
Orius laevigatus, 597
reproduction, 597
sex ratio, 236
biology, 252, 603
armored scale, 252
Asopinae, 603
biological control, 252
giant reed, 252
natural enemy, 603
non-native invasive grass, 252
photophase, 603
stink bug, 603
body size, 750, 775
Coccinellidae, 750
egg size, 750
invasive species, 750
life history strategy, 750
Megachile rotundata, 775
trap-nest, 775
Bombus,
antennal sensilla, 654
rough set, 654
Brazil,
Myrmicinae, 181
Myrmicocrypta bucki sp. nov., 181
Myrmicocrypta camargoi sp. nov., 181
Myrmicocrypta erectapilosa sp. nov., 181
Brevipalpus spp.,
citrus leprosis, 898
SCAR markers, 898
whole genome amplification, 898
brood production,
Dendroctonus pseudotsugae, 39
Dendroctonus simplex, 39
exotic species, 39
Scolytidae, 39
butterfly,
Pieris rapae, 988
sensory bristle, 988
sensory scale, 988
wing margin, 988

C
Castanopsis,
Curculio hilgendorfi, 379
glacial refugia, 379
phylogeography, 379
Rhynchaenus dorsoplanatus, 379
Central, Guanacaste, and Talamanca mountain ranges,
diversity partitioning, 956
light traps, 956
α, β, and γ diversity, 956
Ceratitis capitata, 264, 723
Anastrepha ludens, 264
Anastrepha suspensa, 264
Anastrepha spp., 723
fitness testing, 264
host fruit, 723
long-term cold storage, 264
Neosilba spp., 723
taxonomy, 723
cercus,
character displacement, 327
geographic variation, 327
Parapodisma, 327
cereals,
fitness, 886
mitochondrial DNA, 886
nuclear DNA, 886
sweet corn, 886
character displacement,
cercus, 327
geographic variation, 327
Parapodisma, 327
China, 7, 866
Elateridae, 866
Hypogastruridae, 7
new genera and species, 866
North Dakota, 7
Qinghai Province, 7
species key, 7
Upper Jurassic–Lower Cretaceous, 866
Yixian Formation, 866
cicadas, 216, 860
acoustic behavior, 860
diversity, 216
exuviae, 216
morphometrics, 860
plantation, 216
systematics, 860
thermal adaptation, 860
tropical forest, 216
citrus greening disease,
Diaphorina citri, 611
huanglongbing, 611
rearing, 611
citrus leprosis,
Brevipalpus spp., 898
SCAR markers, 898
whole genome amplification, 898
citrus mealybug,
DC-electrical penetration graph, 404
feeding behavior, 404
grapevine, 404
Planococcus citri, 404
cloning,
Antheraea pernyi, 647
expression pattern, 647
lysophospholipase, 647
Coccinellidae,
body size, 750
egg size, 750
invasive species, 750
life history strategy, 750
codling moth,
ATP, 418
controlled atmospheres, 418
differential scanning calorimetry, 418
metabolism, 418
coexistence, 757, 936
desiccation, 757
Dineulophus phtorimaeae, 936
eggs, 757
extinction, 757
invasive mosquitoes, 757
parasitoid, 936
Pseudapanteles dignus, 936
Tuta absoluta, 936
Coleoptera, 1, 785
conservation genetics, 785
Derolathrus cavernicolus n. sp., 1
island relationships, 785
Jacobsoniidae, 1
multi-island endemic, 785
Tenebrionidae, 785
Collembola,
Appalachian Mountains, 472
phylogeny, 472
taxonomy, 472
Tomoceridae, 472
colony decline,
Acarapis woodi, 711
Apis mellifera, 711
varroa mite, 711
community structure,
biodiversity, 71
biological conservation, 71
faunal survey, 71
functional groups, 71
competition,
container mosquito, 734
density dependence, 734
invasive species, 734
life table, 734
confidence Intervals,
Diaphorina, 827
pooling, 827
prevalence, 827
vector, 827
conservation,
genetic differentiation, 625
genetic diversity, 625
glacial, 625
refugium, 625
conservation genetics, 785, 1012
Coleoptera, 785
dragonfly, 1012
island relationships, 785
multi-island endemic, 785
Somatochlora hineana, 1012
Tenebrionidae, 785
contact pheromones,
cuticular hydrocarbons, 128
Ellychnia corrusca, 128
mating behavior, 128
solid phase microextraction, 128
container community,
antipredatory behavior, 1038
predation risk, 1038
container mosquito,
competition, 734
density dependence, 734
invasive species, 734
life table, 734
controlled atmospheres,
ATP, 418
codling moth, 418
differential scanning calorimetry, 418
metabolism, 418
Cretaceous,
evolution, 875
Jurassic, 875
northeastern China, 875
Pelecinidae, 875
Croatia,
Ucka Nature Park, 165
faunistic survey, 165
Typhlodrominae, 165
Typhlodromus (*Typhlodromus*) *octogenipilus*, 165
crop,
Apis mellifera, 979
cuticle, 979
honey bee, 979
honey stomach, 979
cryptic,
cytotoxicity, 273
hemocyte count, 273
hemocyte viability, 273
mantid, 273
natural enemy, 273
primary defenses, 273
secondary defenses, 273
wasp venom, 273
cryptic species,
armored scale, 844
Diaspididae, 844

oleander scale, 844
species delimitation, 844
Cucurbitaceae,
Aphidiinae, 153
aphids, 153
Solanaceae, 153
southeastern Europe, 153
Culicidae,
Aedes, 618
life history trait, 618
Nepenthes pitcher plant, 618
Stegomyia, 618
Curculio hilgendorfi,
Castanopsis, 379
glacial refugia, 379
phylogeography, 379
Rhynchaenus dorsoplanatus, 379
cuticle,
Apis mellifera, 979
crop, 979
honey bee, 979
honey stomach, 979
cuticular hydrocarbons,
contact pheromones, 128
Ellychnia corrusca, 128
mating behavior, 128
solid phase microextraction, 128
cutworm,
Abagrotis orbis, 592
Agrotis vetusta, 592
grape, 592
Noctuidae, 592
Cynipidae,
inquilines, 492
new genus, 492
Quercus, 492
Synergini, 492
cytochrome oxidase 1,
barcoding, 196
phylogenetics, 196
whitefly, 196
cytochrome oxidase I,
amplified fragment length polymorphism, 1018
genetic diversity, 1018
Troides aeacus, 1018
cytotoxicity, 273, 293
cryptic, 273
hemocyte count, 273, 293
hemocyte viability, 273, 293
mantid, 273
natural enemy, 273
primary defenses, 273
secondary defenses, 273
wasp venom, 273, 293

D

Daohugou,
Gumillinae, 855
Jurassic, 855
Neuroptera, 855
Osmylidae, 855
DC-electrical penetration graph,
citrus mealybug, 404
feeding behavior, 404
grapevine, 404
Planococcus citri, 404
death frequency,
age-specific mortality, 574
Plutella xylostella, 574
defense,
fire ants, 678
nest disturbance, 678
Solenopsis invicta, 678
worker size, 678
Dendroctonus pseudotsugae,
brood production, 39
Dendroctonus simplex, 39
exotic species, 39
Scolytidae, 39
Dendroctonus simplex,
brood production, 39
Dendroctonus pseudotsugae, 39
exotic species, 39
Scolytidae, 39
density dependence,
competition, 734
container mosquito, 734
invasive species, 734
life table, 734
deposition,
excavation, 449
idiothetic, 449
path integration, 449
tunnel, 449
Derolathrus cavernicolus n. sp.,
Coleoptera, 1
Jacobsoniidae, 1
desert,
biogeography, 555
phylogeography, 555
Sphaeropthalminae, 555
velvet ants, 555
desiccation,
coexistence, 757
eggs, 757
extinction, 757
invasive mosquitoes, 757
development,
diet, 66
fecundity, 66
Pezothrips kellyanus, 66
pollen, 66
developmental time,
Planococcus ficus, 943
temperature threshold, 943
thermal requirements, 943
vines, 943
Diabrotica barberi,
fecundity, 688
longevity, 688
mate choice, 688
Diachasma alloeum,
Diachasmimorpha mellea, 908
Rhagoletis pomonella, 908
sequential speciation, 908
Utetes canaliculatus, 908
Diachasmimorpha mellea,
Diachasma alloeum, 908
Rhagoletis pomonella, 908
sequential speciation, 908
Utetes canaliculatus, 908
diapause,
arrest of reproduction, 366
Leptocorisa chinensis, 366
recovery of reproduction, 366
Diaphorina citri,
citrus greening disease, 611
huanglongbing, 611
rearing, 611
Diaphorina,
confidence Intervals, 827
pooling, 827
prevalence, 827
vector, 827
Diaspididae,
armored scale, 844
cryptic species, 844
oleander scale, 844
species delimitation, 844
diet,
development, 66
fecundity, 66
Pezothrips kellyanus, 66
pollen, 66
differential scanning calorimetry,
ATP, 418
codling moth, 418
controlled atmospheres, 418
metabolism, 418
Dineulophus phtorimaeae,
coexistence, 936
parasitoid, 936
Pseudapanteles dignus, 936
Tuta absoluta, 936
Diptera,
Axymyiidae, 455
Jurassic, 455
male adults, 455
taxa, 455
disturbance,
richness, 635
Solenopsis invicta, 635
species composition, 635
vegetation structure, 635
diversity,
cicadas, 216
exuviae, 216
plantation, 216
tropical forest, 216
diversity partitioning,
Central, Guanacaste, and Talamanca mountain ranges, 956

light traps, 956
α, β, and γ diversity, 956
DNA,
invasive species, 548
recent introduction, 548
DNA barcode,
Aethina tumida, 671
Apis mellifera, 671
invasive species, 671
pollination, 671
DNA barcoding,
forest defoliator, 135
hybridization, 135
invasive species, 135
pheromone trap survey, 135
DNA fingerprinting,
Asian citrus psyllid, 100
biological control, 100
ectoparasitoid, 100
natural enemy, 100
dragonfly,
conservation genetics, 1012
Somatochlora hineana, 1012
Drosophila,
fitness, 247
offspring number, 247
perfluorooctane sulfonic acid, 247
terrestrial, 247

E

East and South-East Asia,
Melolontha, 341
Melolonthinae, 341
Scarabaeidae, 341
eastern Palearctic,
alternation of generations, 838
new species, 838
oak-gall wasp, 838
ring-cupped oak, 838
ectoparasitoid,
Asian citrus psyllid, 100
biological control, 100
DNA fingerprinting, 100
natural enemy, 100
egg development,
enzyme-linked immunosorbent assay, 971
Ephestia kuehniella, 971
nutrition, 971
yolk protein, 971
egg elongation,
female body size, 580
grass-feeding butterfly, 580
life-history traits, 580
egg size,
body size, 750
Coccinellidae, 750
invasive species, 750
life history strategy, 750
eggs,
coexistence, 757
desiccation, 757
extinction, 757
invasive mosquitoes, 757
Eichhornia crassipes,
Neotropical, 332
Taosa herbida, 332
Taosa inexacta, 332
Taosa (*Cuernavaca*) *longula* sp. nov., 332
ejaculate expulsion,
natural history, 111
Ulidiidae, 111
Elachipterini,
Africa, 465
Oscinellinae, 465
phylogeny, 465
taxonomy, 465
Elateridae,
China, 866
new genera and species, 866
Upper Jurassic–Lower Cretaceous, 866
Yixian Formation, 866
Ellychnia corrusca,
contact pheromones, 128
cuticular hydrocarbons, 128
mating behavior, 128
solid phase microextraction, 128
entomopathogenic nematodes,
alkaloids, 371
pathogen reproduction, 371
polyphagy, 371
trade-off, 371
enzyme-linked immunosorbent assay,
egg development, 971
Ephestia kuehniella, 971
nutrition, 971
yolk protein, 971
Ephestia kuehniella,
egg development, 971
enzyme-linked immunosorbent assay, 971
nutrition, 971
yolk protein, 971
eriophyoid mites,
key, 697
new species, 697
plant feeding, 697
taxonomy, 697
evolution,
Cretaceous, 875
Jurassic, 875
northeastern China, 875
Pelecinidae, 875
excavation,
deposition, 449
idiothetic, 449
path integration, 449
tunnel, 449
exotic species,
brood production, 39
Dendroctonus pseudotsugae, 39
Dendroctonus simplex, 39
Scolytidae, 39
expression pattern,
Antheraea pernyi, 647
cloning, 647
lysophospholipase, 647
extinction,
coexistence, 757
desiccation, 757
eggs, 757
invasive mosquitoes, 757
exuviae,
cicadas, 216
diversity, 216
plantation, 216
tropical forest, 216

F

faunal survey,
biodiversity, 71
biological conservation, 71
community structure, 71
functional groups, 71
faunistic survey,
Ucka Nature Park, 165
Croatia, 165
Typhlodrominae, 165
Typhlodromus (*Typhlodromus*) *octogenipilus*, 165
fecundity, 57, 66, 397, 688
Ascidae, 397
development, 66
Diabrotica barberi, 688
diet, 66
flower mites, 397
glassy-winged sharpshooter, 57
leafhopper, 57
longevity, 57, 688
mate choice, 688
Pezothrips kellyanus, 66
pollen, 66
pollen feeding, 397
rearing density, 57
feeding behavior, 404, 585
citrus mealybug, 404
DC-electrical penetration graph, 404
grapevine, 404
Heteroptera, 585
host plant, 585
Planococcus citri, 404
Poaceae, 585
female body size,
egg elongation, 580
grass-feeding butterfly, 580
life-history traits, 580

female mating receptivity,
age, 802
male courtship, 802
Miridae, 802
ovarian development, 802
fire ants,
defense, 678
nest disturbance, 678
Solenopsis invicta, 678
worker size, 678
first-feeding delay,
locomotion performance, 439
oviposition on nonhosts, 439
relative leg length, 439
synthetic aggregation pheromone, 439
fitness, 247, 886
cereals, 886
Drosophila, 247
mitochondrial DNA, 886
nuclear DNA, 886
offspring number, 247
perfluorooctane sulfonic acid, 247
sweet corn, 886
terrestrial, 247
fitness testing,
Anastrepha ludens, 264
Anastrepha suspensa, 264
Ceratitis capitata, 264
long-term cold storage, 264
Florida,
behavioral choice, 227
local adaptation, 227
oviposition choice, 227
phenotypic variation, 227
flower mites,
Ascidae, 397
fecundity, 397
pollen feeding, 397
fluorescence in situ hybridization,
Hymenoptera, 424
karyotype, 424
Ponerinae, 424
forest defoliator,
DNA barcoding, 135
hybridization, 135
invasive species, 135
pheromone trap survey, 135
forest fires,
infrared radiation sensors, 823
Melanophila acuminata, 823
reproduction, 823
Formicidae,
morphometrics, 20
multivariate allometry, 20
Tapinoma schreiberi, 20
Tapinoma sessile, 20
functional groups,
biodiversity, 71
biological conservation, 71
community structure, 71
faunal survey, 71
fungus weevils,
Anthribinae, 523
morphology, 523
phylogeny, 523
Ptychoderini, 523

G

generic description,
Alydinae, 706
Leptocorisini, 706
Melanacanthus, 706
Micrelytrinae, 706
genetic differentiation, 209, 625
allometry, 209
conservation, 625
genetic diversity, 625
glacial, 625
microevolution, 209
phenotypic plasticity, 209
refugium, 625
sexual selection, 209
genetic diversity, 625, 1018
amplified fragment length polymorphism, 1018
conservation, 625
cytochrome oxidase I, 1018
genetic differentiation, 625
glacial, 625
refugium, 625
Troides aeacus, 1018
geographic variation,
cercus, 327
character displacement, 327
Parapodisma, 327
geometric morphometry,
Apis, 965
honey bee, 965
wing morphometry, 965
giant reed,
armored scale, 252
biological control, 252
biology, 252
non-native invasive grass, 252
glacial,
conservation, 625
genetic differentiation, 625
genetic diversity, 625
refugium, 625
glacial refugia,
Castanopsis, 379
Curculio hilgendorfi, 379
phylogeography, 379
Rhynchaenus dorsoplanatus, 379
glassy-winged sharpshooter, 57, 236
biological control, 236
fecundity, 57
Homalodisca liturata, 236
leafhopper, 57
longevity, 57
morphometrics, 236
rearing density, 57
sex ratio, 236
grape,
Abagrotis orbis, 592
Agrotis vetusta, 592
cutworm, 592
Noctuidae, 592
grapevine,
citrus mealybug, 404
DC-electrical penetration graph, 404
feeding behavior, 404
Planococcus citri, 404
grass-feeding butterfly, 580, 716
egg elongation, 580
female body size, 580
life-history strategy, 716
life-history traits, 580
mitochondrial DNA, 716
phylogenetic constraints, 716
seasonal adaptation, 716
Greenideinae,
Aphidiinae, 307
key, 307
review, 307
Gumillinae,
Daohugou, 855
Jurassic, 855
Neuroptera, 855
Osmylidae, 855

H

haplotype diversity,
intraspecific variation, 916
nucleotide diversity, 916
phylogenetic tree, 916
Helotidae,
Laos, 500
Myanmar, 500
new species, 500
taxonomic revision, 500
Hemiptera, 544, 813
adaptation of populations, 544
leafhoppers, 544
maize, 544
phytoplasma vector, 813
planthopper, 813
vibrational signals, 813
hemocyte count, 273, 293
cryptic, 273
cytotoxicity, 273, 293
hemocyte viability, 273, 293
mantid, 273
natural enemy, 273
primary defenses, 273
secondary defenses, 273
wasp venom, 273, 293
hemocyte viability, 273, 293
cryptic, 273
cytotoxicity, 273, 293
hemocyte count, 273, 293

mantid, 273
natural enemy, 273
primary defenses, 273
secondary defenses, 273
wasp venom, 273, 293
heterogony,
Andricus, 322
life cycle, 322
synonym, 322
Heteroptera,
feeding behavior, 585
host plant, 585
Poaceae, 585
Homalodisca liturata,
biological control, 236
glassy-winged sharpshooter, 236
morphometrics, 236
sex ratio, 236
honey bee, 146, 389, 965, 979
Aethina tumida, 389
Apis mellifera, 979
Apis, 965
crop, 979
cuticle, 979
geometric morphometry, 965
honey stomach, 979
hygiene, 146
Kenya, 389
Kodamaea ohmeri, 389
Oplostomus haroldi, 389
varroa, 146
varroa sensitive hygiene, 146
wing morphometry, 965
honey stomach,
Apis mellifera, 979
crop, 979
cuticle, 979
honey bee, 979
host alternation,
Aphis glycines, 532
Aphis gossypii, 532
molecular identification, 532
morphology, 532
host fruit,
Anastrepha spp., 723
Ceratitis capitata, 723
Neosilba spp., 723
taxonomy, 723
host plant,
feeding behavior, 585
Heteroptera, 585
Poaceae, 585
Host race,
incomplete lineage sorting, 11
mitochondrial DNA capture, 11
phytophagous insects, 11
host strains,
Spodoptera frugiperda, 283
triose phosphate isomerase, 283
huanglongbing,
citrus greening disease, 611
Diaphorina citri, 611
rearing, 611
hybridization,
DNA barcoding, 135
forest defoliator, 135
invasive species, 135
pheromone trap survey, 135
hygiene,
honey bees, 146
varroa, 146
varroa sensitive hygiene, 146
Hymenoptera,
fluorescence in situ hybridization, 424
karyotype, 424
Ponerinae, 424
Hypogastruridae,
China, 7
North Dakota, 7
Qinghai Province, 7
species key, 7

I

idiothetic,
deposition, 449
excavation, 449
path integration, 449
tunnel, 449
imidacloprid,
metabolism, 84
Reticulitermes flavipes, 84
termite, 84
toxicity, 84
incomplete lineage sorting,
Host race, 11
mitochondrial DNA capture, 11
phytophagous insects, 11
infrared radiation sensors,
forest fires, 823
Melanophila acuminata, 823
reproduction, 823
inheritance,
Anastrepha ludens, 96
linkage analysis, 96
mutations, 96
inquilines,
Cynipidae, 492
new genus, 492
Quercus, 492
Synergini, 492
insect behavior,
parental investment, 771
Zelus, 771
insect pest,
Abgrallaspis, 30
armored scale insects, 30
avocado, 30
molecular phylogenetics, 30
internal transcribed spacer,
Bayesian inference, 742
mitochondrial DNA, 742
parasitoids, 742
spatial genetic structure, 742
intraspecific variation,
haplotype diversity, 916
nucleotide diversity, 916
phylogenetic tree, 916
introgression,
life-history traits, 925
northern corn rootworm, 925
population growth rates, 925
invasive mosquitoes,
coexistence, 757
desiccation, 757
eggs, 757
extinction, 757
invasive species, 135, 548, 671, 734, 750
Aethina tumida, 671
Apis mellifera, 671
body size, 750
Coccinellidae, 750
competition, 734
container mosquito, 734
density dependence, 734
DNA, 548
DNA barcode, 671
DNA barcoding, 135
egg size, 750
forest defoliator, 135
hybridization, 135
life history strategy, 750
life table, 734
pheromone trap survey, 135
pollination, 671
recent introduction, 548
island relationships,
Coleoptera, 785
conservation genetics, 785
multi-island endemic, 785
Tenebrionidae, 785
Ithonidae,
lacewing, 511
phylogeny, 511
Polystoechotidae, 511
Rapismatidae, 511

J

Jacobsoniidae,
Coleoptera, 1
Derolathrus cavernicolus n. sp., 1
Jurassic, 455, 855, 875
Axymyiidae, 455
Cretaceous, 875
Daohugou, 855
Diptera, 455
evolution, 875
Gumillinae, 855
male adults, 455

Neuroptera, 855
northeastern China, 875
Osmylidae, 855
Pelecinidae, 875
taxa, 455

K

karyotype,
fluorescence in situ hybridization, 424
Hymenoptera, 424
Ponerinae, 424
Kenya,
Aethina tumida, 389
honey bee, 389
Kodamaea ohmeri, 389
Oplostomus haroldi, 389
key, 307, 697
Aphidiinae, 307
eriophyoid mites, 697
Greenideinae, 307
new species, 697
plant feeding, 697
review, 307
taxonomy, 697
Kodamaea ohmeri,
Aethina tumida, 389
honey bee, 389
Kenya, 389
Oplostomus haroldi, 389

L

lacewing,
Ithonidae, 511
phylogeny, 511
Polystoechotidae, 511
Rapismatidae, 511
Laos,
Helotidae, 500
Myanmar, 500
new species, 500
taxonomic revision, 500
leaf anatomy,
leaf nutrition, 806
leafminer, 806
mining pattern, 806
Ophiomyia maura, 806
leaf nutrition,
leaf anatomy, 806
leafminer, 806
mining pattern, 806
Ophiomyia maura, 806
leafhopper, 57, 544
adaptation of populations, 544
fecundity, 57
glassy-winged sharpshooter, 57
Hemiptera, 544
longevity, 57
maize, 544
rearing density, 57
leafminer,
leaf anatomy, 806
leaf nutrition, 806
mining pattern, 806
Ophiomyia maura, 806
Leptocorisa chinensis,
arrest of reproduction, 366
diapause, 366
recovery of reproduction, 366
Leptocorisini,
Alydinae, 706
generic description, 706
Melanacanthus, 706
Micrelytrinae, 706
lesser grain borer,
aggregation pheromone, 796
beetle density, 796
nutritional stress, 796
pheromone output, 796
life cycle,
Andricus, 322
heterogony, 322
synonym, 322
life history strategy,
body size, 750
Coccinellidae, 750
egg size, 750
invasive species, 750
life history trait,
Aedes, 618
Culicidae, 618
Nepenthes pitcher plant, 618
Stegomyia, 618
life table,
competition, 734
container mosquito, 734
density dependence, 734
invasive species, 734
life-history strategy,
grass-feeding butterfly, 716
mitochondrial DNA, 716
phylogenetic constraints, 716
seasonal adaptation, 716
life-history traits, 580, 925
egg elongation, 580
female body size, 580
grass-feeding butterfly, 580
introgression, 925
northern corn rootworm, 925
population growth rates, 925
light traps,
Central, Guanacaste, and Talamanca mountain ranges, 956
diversity partitioning, 956
α, β, and γ diversity, 956
linkage analysis,
Anastrepha ludens, 96
inheritance, 96
mutations, 96
local adaptation,
behavioral choice, 227
Florida, 227
oviposition choice, 227
phenotypic variation, 227
locomotion performance,
first-feeding delay, 439
oviposition on nonhosts, 439
relative leg length, 439
synthetic aggregation pheromone, 439
long-term cold storage,
Anastrepha ludens, 264
Anastrepha suspensa, 264
Ceratitis capitata, 264
fitness testing, 264
longevity, 57, 688
Diabrotica barberi, 688
fecundity, 57, 688
glassy-winged sharpshooter, 57
leafhopper, 57
mate choice, 688
rearing density, 57
Lygus hesperus,
postmating refractory period, 300
sexual receptivity, 300
spermatophore, 300
Lymantria monacha,
artificial diet, 949
phenology, 949
rearing, 949
lysophospholipase,
Antheraea pernyi, 647
cloning, 647
expression pattern, 647

M

maize,
adaptation of populations, 544
Hemiptera, 544
leafhoppers, 544
male adults,
Axymyiidae, 455
Diptera, 455
Jurassic, 455
taxa, 455
male courtship,
age, 802
female mating receptivity, 802
Miridae, 802
ovarian development, 802
mantid, 120, 273
cryptic, 273
cytotoxicity, 273
hemocyte count, 273
hemocyte viability, 273
Mantodea, 120
natural enemy, 273
pheromones, 120
primary defenses, 273
secondary defenses, 273

Stagmomantis limbata, 120
Tenodera aridifolia sinensis, 120
wasp venom, 273
Mantodea,
mantid, 120
pheromones, 120
Stagmomantis limbata, 120
Tenodera aridifolia sinensis, 120
mate choice,
Diabrotica barberi, 688
fecundity, 688
longevity, 688
mating,
artificial diet, 597
biological control, 597
Orius laevigatus, 597
reproduction, 597
mating behavior, 128, 430
agave, 430
contact pheromones, 128
cuticular hydrocarbons, 128
Ellychnia corrusca, 128
natural history, 430
solid phase microextraction, 128
Tephritoidea, 430
Ulidiidae, 430
Megachile rotundata,
body size, 775
trap-nest, 775
Melanacanthus,
Alydinae, 706
generic description, 706
Leptocorisini, 706
Micrelytrinae, 706
Melanophila acuminata,
forest fires, 823
infrared radiation sensors, 823
reproduction, 823
Melolontha,
East and South-East Asia, 341
Melolonthinae, 341
Scarabaeidae, 341
Melolonthinae,
East and South-East Asia, 341
Melolontha, 341
Scarabaeidae, 341
metabolism, 84, 418
ATP, 418
codling moth, 418
controlled atmospheres, 418
differential scanning calorimetry, 418
imidacloprid, 84
Reticulitermes flavipes, 84
termite, 84
toxicity, 84
Micrelytrinae,
Alydinae, 706
generic description, 706
Leptocorisini, 706
Melanacanthus, 706
microevolution,
allometry, 209
genetic differentiation, 209
phenotypic plasticity, 209
sexual selection, 209
Minangini,
biogeography, 360
Oriental, 360
Palearctic, 360
Sigalphinae, 360
mining pattern,
leaf anatomy, 806
leaf nutrition, 806
leafminer, 806
Ophiomyia maura, 806
Miridae,
age, 802
female mating receptivity, 802
male courtship, 802
ovarian development, 802
mitochondrial DNA, 716, 742, 886, 1025
Bayesian inference, 742
cereals, 886
fitness, 886
grass-feeding butterfly, 716
internal transcribed spacer, 742
life-history strategy, 716
nuclear DNA, 886
parasitoids, 742
phylogenetic constraints, 716
phylogeography, 1025
seasonal adaptation, 716
spatial genetic structure, 742
sweet corn, 886
Wallacea, 1025
mitochondrial DNA capture,
Host race, 11
incomplete lineage sorting, 11
phytophagous insects, 11
mitotic chromosomes,
agricultural pest, 1003
polytene chromosomes, 1003
Rhagoletis completa, 1003
Wolbachia pipientis, 1003
molecular identification,
Aphis glycines, 532
Aphis gossypii, 532
host alternation, 532
morphology, 532
molecular phylogenetics,
Abgrallaspis, 30
armored scale insects, 30
avocado, 30
insect pest, 30
morphology, 523, 532
Anthribinae, 523
Aphis glycines, 532
Aphis gossypii, 532
fungus weevils, 523
host alternation, 532
molecular identification, 532
phylogeny, 523
Ptychoderini, 523
morphometrics, 20, 236, 860
acoustic behavior, 860
biological control, 236
cicadas, 860
Formicidae, 20
glassy-winged sharpshooter, 236
Homalodisca liturata, 236
multivariate allometry, 20
sex ratio, 236
systematics, 860
Tapinoma schreiberi, 20
Tapinoma sessile, 20
thermal adaptation, 860
mouthpart length,
Asteraceae, 566
nectar feeding, 566
parasitic wasps, 566
multi-island endemic,
Coleoptera, 785
conservation genetics, 785
island relationships, 785
Tenebrionidae, 785
multivariate allometry,
Formicidae, 20
morphometrics, 20
Tapinoma schreiberi, 20
Tapinoma sessile, 20
mutations,
Anastrepha ludens, 96
inheritance, 96
linkage analysis, 96
Myanmar,
Helotidae, 500
Laos, 500
new species, 500
taxonomic revision, 500
Myrmicinae,
Brazil, 181
Myrmicocrypta bucki sp. nov., 181
Myrmicocrypta camargoi sp. nov., 181
Myrmicocrypta erectapilosa sp. nov., 181
Myrmicocrypta bucki sp. nov.,
Brazil, 181
Myrmicinae, 181
Myrmicocrypta camargoi sp. nov., 181
Myrmicocrypta erectapilosa sp. nov., 181
Myrmicocrypta camargoi sp. nov.,
Brazil, 181
Myrmicinae, 181
Myrmicocrypta bucki sp. nov., 181

Myrmicocrypta erectapilosa sp. nov., 181
Myrmicocrypta erectapilosa sp. nov., Brazil, 181
Myrmicinae, 181
Myrmicocrypta bucki sp. nov., 181
Myrmicocrypta camargoi sp. nov., 181

N

natural enemy, 100, 273, 603
Asian citrus psyllid, 100
Asopinae, 603
biological control, 100
biology, 603
cryptic, 273
cytotoxicity, 273
DNA fingerprinting, 100
ectoparasitoid, 100
hemocyte count, 273
hemocyte viability, 273
mantid, 273
photophase, 603
primary defenses, 273
secondary defenses, 273
stink bug, 603
wasp venom, 273
natural history, 111, 430
agave, 430
ejaculate expulsion, 111
mating behavior, 430
Tephritoidea, 430
Ulidiidae, 111, 430
nectar feeding,
Asteraceae, 566
mouthpart length, 566
parasitic wasps, 566
negative geotaxis,
Sitophilus oryzae, 683
Sitophilus zeamais, 683
stored grain insects, 683
Neosilba spp.,
Anastrepha spp., 723
Ceratitis capitata, 723
host fruit, 723
taxonomy, 723
Neotropical,
Eichhornia crassipes, 332
Taosa herbida, 332
Taosa inexacta, 332
Taosa (*Cuernavaca*) *longula* sp. nov., 332
Nepenthes pitcher plant,
Aedes, 618
Culicidae, 618
life history trait, 618
Stegomyia, 618
nest disturbance,
defense, 678
fire ants, 678
Solenopsis invicta, 678
worker size, 678
Neuroptera,
Daohugou, 855
Gumillinae, 855
Jurassic, 855
Osmylidae, 855
new genera and species,
China, 866
Elateridae, 866
Upper Jurassic–Lower Cretaceous, 866
Yixian Formation, 866
new genus,
Cynipidae, 492
inquilines, 492
Quercus, 492
Synergini, 492
new species, 500, 697, 838
alternation of generations, 838
eastern Palearctic, 838
eriophyoid mites, 697
Helotidae, 500
key, 697
Laos, 500
Myanmar, 500
oak-gall wasp, 838
plant feeding, 697
ring-cupped oak, 838
taxonomic revision, 500
taxonomy, 697
Noctuidae,
Abagrotis orbis, 592
Agrotis vetusta, 592
cutworm, 592
grape, 592
non-native invasive grass,
armored scale, 252
biological control, 252
biolc,;y, 252
giant reed, 252
North Dakota,
China, 7
Hypogastruridae, 7
Qinghai Province, 7
species key, 7
northeastern China,
Cretaceous, 875
evolution, 875
Jurassic, 875
Pelecinidae, 875
northern corn rootworm,
introgression, 925
life-history traits, 925
population growth rates, 925
nuclear DNA,
cereals, 886
fitness, 886
mitochondrial DNA, 886
sweet corn, 886
nucleotide diversity,
haplotype diversity, 916
intraspecific variation, 916
phylogenetic tree, 916
nutrition,
egg development, 971
enzyme-linked immunosorbent assay, 971
Ephestia kuehniella, 971
yolk protein, 971
nutritional stress,
aggregation pheromone, 796
beetle density, 796
lesser grain borer, 796
pheromone output, 796

O

oak-gall wasp,
alternation of generations, 838
eastern Palearctic, 838
new species, 838
ring-cupped oak, 838
offspring number,
Drosophila, 247
fitness, 247
perfluorooctane sulfonic acid, 247
terrestrial, 247
oleander scale,
armored scale, 844
cryptic species, 844
Diaspididae, 844
species delimitation, 844
olfaction,
antennal morphology, 993
sensilla, 993
Synanthedon scitula, 993
Ophiomyia maura,
leaf anatomy, 806
leaf nutrition, 806
leafminer, 806
mining pattern, 806
Oplostomus haroldi,
Aethina tumida, 389
honey bee, 389
Kenya, 389
Kodamaea ohmeri, 389
Oriental,
biogeography, 360
Minangini, 360
Palearctic, 360
Sigalphinae, 360
Orius laevigatus,
artificial diet, 597
biological control, 597
mating, 597
reproduction, 597
Oscinellinae,
Africa, 465
Elachipterini, 465
phylogeny, 465
taxonomy, 465
Osmylidae,
Daohugou, 855
Gumillinae, 855

Jurassic, 855
Neuroptera, 855
ovarian development,
age, 802
female mating receptivity, 802
male courtship, 802
Miridae, 802
oviposition choice,
behavioral choice, 227
Florida, 227
local adaptation, 227
phenotypic variation, 227
oviposition on nonhosts,
first-feeding delay, 439
locomotion performance, 439
relative leg length, 439
synthetic aggregation pheromone, 439

P

Palearctic,
biogeography, 360
Minangini, 360
Oriental, 360
Sigalphinae, 360
Parapodisma,
cercus, 327
character displacement, 327
geographic variation, 327
parasitic wasps,
Asteraceae, 566
mouthpart length, 566
nectar feeding, 566
parasitoid, 742, 939
Bayesian inference, 742
coexistence, 936
Dineulophus phtorimaeae, 936
internal transcribed spacer, 742
mitochondrial DNA, 742
Pseudapanteles dignus, 936
spatial genetic structure, 742
Tuta absoluta, 936
parental investment,
insect behavior, 771
Zelus, 771
path integration,
deposition, 449
excavation, 449
idiothetic, 449
tunnel, 449
pathogen reproduction,
alkaloids, 371
entomopathogenic nematodes, 371
polyphagy, 371
trade-off, 371
Pelecinidae,
Cretaceous, 875
evolution, 875
Jurassic, 875
northeastern China, 875
perfluorooctane sulfonic acid,
Drosophila, 247
fitness, 247
offspring number, 247
terrestrial, 247
Pezothrips kellyanus,
development, 66
diet, 66
fecundity, 66
pollen, 66
phenology,
artificial diet, 949
Lymantria monacha, 949
rearing, 949
phenotypic plasticity,
allometry, 209
genetic differentiation, 209
microevolution, 209
sexual selection, 209
phenotypic variation,
behavioral choice, 227
Florida, 227
local adaptation, 227
oviposition choice, 227
pheromone output,
aggregation pheromone, 796
beetle density, 796
lesser grain borer, 796
nutritional stress, 796
pheromone trap survey,
DNA barcoding, 135
forest defoliator, 135
hybridization, 135
invasive species, 135
pheromones,
mantid, 120
Mantodea, 120
Stagmomantis limbata, 120
Tenodera aridifolia sinensis, 120
photophase,
Asopinae, 603
biology, 603
natural enemy, 603
stink bug, 603
phylogenetic constraints,
grass-feeding butterfly, 716
life-history strategy, 716
mitochondrial DNA, 716
seasonal adaptation, 716
phylogenetic tree,
haplotype diversity, 916
intraspecific variation, 916
nucleotide diversity, 916
phylogenetics,
barcoding, 196
cytochrome oxidase 1, 196
whitefly, 196
phylogeny, 465, 472, 511, 523
Africa, 465
Anthribinae, 523
Appalachian Mountains, 472
Collembola, 472
Elachipterini, 465
fungus weevils, 523
Ithonidae, 511
lacewing, 511
morphology, 523
Oscinellinae, 465
Polystoechotidae, 511
Ptychoderini, 523
Rapismatidae, 511
taxonomy, 465, 472
Tomoceridae, 472
phylogeography, 379, 555, 1025
biogeography, 555
Castanopsis, 379
Curculio hilgendorfi, 379
desert, 555
glacial refugia, 379
mitochondrial DNA, 1025
Rhynchaenus dorsoplanatus, 379
Sphaeropthalminae, 555
velvet ants, 555
Wallacea, 1025
phytophagous insects,
Host race, 11
incomplete lineage sorting, 11
mitochondrial DNA capture, 11
phytoplasma vector,
Hemiptera, 813
planthopper, 813
vibrational signals, 813
Pieris rapae,
butterfly, 988
sensory bristle, 988
sensory scale, 988
wing margin, 988
Planococcus citri,
citrus mealybug, 404
DC-electrical penetration graph, 404
feeding behavior, 404
grapevine, 404
Planococcus ficus,
developmental time, 943
temperature threshold, 943
thermal requirements, 943
vines, 943
plant feeding,
eriophyoid mites, 697
key, 697
new species, 697
taxonomy, 697
plantation,
cicadas, 216
diversity, 216
exuviae, 216
tropical forest, 216
planthopper,
Hemiptera, 813
phytoplasma vector, 813

vibrational signals, 813
Plutella xylostella,
age-specific mortality, 574
death frequency, 574
Poaceae,
feeding behavior, 585
Heteroptera, 585
host plant, 585
pollen,
development, 66
diet, 66
fecundity, 66
Pezothrips kellyanus, 66
pollen feeding,
Ascidae, 397
fecundity, 397
flower mites, 397
pollination,
Aethina tumida, 671
Apis mellifera, 671
DNA barcode, 671
invasive species, 671
polyphagy,
alkaloids, 371
entomopathogenic nematodes, 371
pathogen reproduction, 371
trade-off, 371
Polystoechotidae,
Ithonidae, 511
lacewing, 511
phylogeny, 511
Rapismatidae, 511
polytene chromosomes,
agricultural pest, 1003
mitotic chromosomes, 1003
Rhagoletis completa, 1003
Wolbachia pipientis, 1003
Ponerinae,
fluorescence in situ hybridization, 424
Hymenoptera, 424
karyotype, 424
pooling,
confidence Intervals, 827
Diaphorina, 827
prevalence, 827
vector, 827
population growth rates,
introgression, 925
life-history traits, 925
northern corn rootworm, 925
postmating refractory period,
Lygus hesperus, 300
sexual receptivity, 300
spermatophore, 300
predation risk,
antipredatory behavior, 1038
container community, 1038
prevalence,
confidence Intervals, 827
Diaphorina, 827
pooling, 827
vector, 827
primary defenses,
cryptic, 273
cytotoxicity, 273
hemocyte count, 273
hemocyte viability, 273
mantid, 273
natural enemy, 273
secondary defenses, 273
wasp venom, 273
Pseudapanteles dignus,
coexistence, 936
Dineulophus phtorimaeae, 936
parasitoid, 936
Tuta absoluta, 936
Ptychoderini,
Anthribinae, 523
fungus weevils, 523
morphology, 523
phylogeny, 523

Q

Qinghai Province,
China, 7
Hypogastruridae, 7
North Dakota, 7
species key, 7
Quercus,
Cynipidae, 492
inquilines, 492
new genus, 492
Synergini, 492

R

Rapismatidae,
Ithonidae, 511
lacewing, 511
phylogeny, 511
Polystoechotidae, 511
rearing, 50, 611, 949
artificial diet, 949
citrus greening disease, 611
Diaphorina citri, 611
huanglongbing, 611
Lymantria monacha, 949
phenology, 949
Synanthedon scitula, 50
thinning apples, 50
rearing density,
fecundity, 57
glassy-winged sharpshooter, 57
leafhopper, 57
longevity, 57
recent introduction,
DNA, 548
invasive species, 548
recovery of reproduction,
arrest of reproduction, 366
diapause, 366
Leptocorisa chinensis, 366
refugium,
conservation, 625
genetic differentiation, 625
genetic diversity, 625
glacial, 625
relative leg length,
first-feeding delay, 439
locomotion performance, 439
oviposition on nonhosts, 439
synthetic aggregation pheromone, 439
reproduction, 597, 823
artificial diet, 597
biological control, 597
forest fires, 823
infrared radiation sensors, 823
mating, 597
Melanophila acuminata, 823
Orius laevigatus, 597
Reticulitermes flavipes,
imidacloprid, 84
metabolism, 84
termite, 84
toxicity, 84
review,
Aphidiinae, 307
Greenideinae, 307
key, 307
Rhagoletis completa,
agricultural pest, 1003
mitotic chromosomes, 1003
polytene chromosomes, 1003
Wolbachia pipientis, 1003
Rhagoletis pomonella,
Diachasma alloeum, 908
Diachasmimorpha mellea, 908
sequential speciation, 908
Utetes canaliculatus, 908
Rhynchaenus dorsoplanatus,
Castanopsis, 379
Curculio hilgendorfi, 379
glacial refugia, 379
phylogeography, 379
richness,
disturbance, 635
Solenopsis invicta, 635
species composition, 635
vegetation structure, 635
ring-cupped oak,
alternation of generations, 838
eastern Palearctic, 838
new species, 838
oak-gall wasp, 838
rough set,
antennal sensilla, 654
Bombus, 654

S

scanning electron microscopy,
antennal morphology, 661
sensilla, 661
Tephritidae, 661

SCAR markers,
Brevipalpus spp, 898
citrus leprosis, 898
whole genome amplification, 898
Scarabaeidae,
East and South-East Asia, 341
Melolontha, 341
Melolonthinae, 341
Scolytidae,
brood production, 39
Dendroctonus pseudotsugae, 39
Dendroctonus simplex, 39
exotic species, 39
seasonal adaptation,
grass-feeding butterfly, 716
life-history strategy, 716
mitochondrial DNA, 716
phylogenetic constraints, 716
secondary defenses,
cryptic, 273
cytotoxicity, 273
hemocyte count, 273
hemocyte viability, 273
mantid, 273
natural enemy, 273
primary defenses, 273
wasp venom, 273
sensilla, 661, 993
antennal morphology, 661, 993
olfaction, 993
scanning electron microscopy, 661
Synanthedon scitula, 993
Tephritidae, 661
sensory bristle,
butterfly, 988
Pieris rapae, 988
sensory scale, 988
wing margin, 988
sensory scale,
butterfly, 988
Pieris rapae, 988
sensory bristle, 988
wing margin, 988
sequential speciation,
Diachasma alloeum, 908
Diachasmimorpha mellea, 908
Rhagoletis pomonella, 908
Utetes canaliculatus, 908
sex ratio,
biological control, 236
glassy-winged sharpshooter, 236
Homalodisca liturata, 236
morphometrics, 236
sexual receptivity,
Lygus hesperus, 300
postmating refractory period, 300
spermatophore, 300
sexual selection,
allometry, 209
genetic differentiation, 209
microevolution, 209
phenotypic plasticity, 209
Sigalphinae,
biogeography, 360
Minangini, 360
Oriental, 360
Palearctic, 360
Sitophilus oryzae,
negative geotaxis, 683
Sitophilus zeamais, 683
stored grain insects, 683
Sitophilus zeamais,
negative geotaxis, 683
Sitophilus oryzae, 683
stored grain insects, 683
Solanaceae,
Aphidiinae, 153
aphids, 153
Cucurbitaceae, 153
southeastern Europe, 153
Solenopsis invicta, 635, 678
defense, 678
disturbance, 635
fire ants, 678
nest disturbance, 678
richness, 635
species composition, 635
vegetation structure, 635
worker size, 678
solid phase microextraction,
contact pheromones, 128
cuticular hydrocarbons, 128
Ellychnia corrusca, 128
mating behavior, 128
Somatochlora hineana,
conservation genetics, 1012
dragonfly, 1012
southeastern Europe,
Aphidiinae, 153
aphids, 153
Cucurbitaceae, 153
Solanaceae, 153
spatial genetic structure,
Bayesian inference, 742
internal transcribed spacer, 742
mitochondrial DNA, 742
parasitoids, 742
species composition,
disturbance, 635
richness, 635
Solenopsis invicta, 635
vegetation structure, 635
species delimitation,
armored scale, 844
cryptic species, 844
Diaspididae, 844
oleander scale, 844
species key,
China, 7
Hypogastruridae, 7
North Dakota, 7
Qinghai Province, 7
spermatophore,
Lygus hesperus, 300
postmating refractory period, 300
sexual receptivity, 300
Sphaeropthalminae,
biogeography, 555
desert, 555
phylogeography, 555
velvet ants, 555
Spodoptera frugiperda,
host strains, 283
triose phosphate isomerase, 283
Stagmomantis limbata,
mantid, 120
Mantodea, 120
pheromones, 120
Tenodera aridifolia sinensis, 120
Stegomyia,
Aedes, 618
Culicidae, 618
life history trait, 618
Nepenthes pitcher plant, 618
stink bug,
Asopinae, 603
biology, 603
natural enemy, 603
photophase, 603
stored grain insects,
negative geotaxis, 683
Sitophilus oryzae, 683
Sitophilus zeamais, 683
sweet corn,
cereals, 886
fitness, 886
mitochondrial DNA, 886
nuclear DNA, 886
Synanthedon scitula, 50, 993
antennal morphology, 993
olfaction, 993
rearing, 50
sensilla, 993
thinning apples, 50
Synergini,
Cynipidae, 492
inquilines, 492
new genus, 492
Quercus, 492
synonym,
Andricus, 322
heterogony, 322
life cycle, 322

synthetic aggregation pheromone,
first-feeding delay, 439
locomotion performance, 439
oviposition on nonhosts, 439
relative leg length, 439
systematics,
acoustic behavior, 860
cicadas, 860
morphometrics, 860
thermal adaptation, 860

T

Taosa herbida,
Eichhornia crassipes, 332
Neotropical, 332
Taosa inexacta, 332
Taosa (*Cuernavaca*) *longula* sp. nov., 332
Taosa inexacta,
Eichhornia crassipes, 332
Neotropical, 332
Taosa herbida, 332
Taosa (*Cuernavaca*) *longula* sp. nov., 332
Taosa (*Cuernavaca*) *longula* sp. nov.,
Eichhornia crassipes, 332
Neotropical, 332
Taosa herbida, 332
Taosa inexacta, 332
Taosa (*Cuernavaca*) *longula* sp. nov., 332
Tapinoma schreiberi,
Formicidae, 20
morphometrics, 20
multivariate allometry, 20
Tapinoma sessile, 20
Tapinoma sessile,
Formicidae, 20
morphometrics, 20
multivariate allometry, 20
Tapinoma schreiberi, 20
taxa,
Axymyiidae, 455
Diptera, 455
Jurassic, 455
male adults, 455
taxonomic revision,
Helotidae, 500
Laos, 500
Myanmar, 500
new species, 500
taxonomy, 465, 472, 697, 723
Africa, 465
Anastrepha spp., 723
Appalachian Mountains, 472
Ceratitis capitata, 723
Collembola, 472
Elachipterini, 465
eriophyoid mites, 697
host fruit, 723
key, 697
Neosilba spp., 723
new species, 697
Oscinellinae, 465
phylogeny, 465, 472
plant feeding, 697
Tomoceridae, 472
temperature threshold,
developmental time, 943
Planococcus ficus, 943
thermal requirements, 943
vines, 943
Tenebrionidae,
Coleoptera, 785
conservation genetics, 785
island relationships, 785
multi-island endemic, 785
Tenodera aridifolia sinensis,
mantid, 120
Mantodea, 120
pheromones, 120
Stagmomantis limbata, 120
Tephritidae,
antennal morphology, 661
scanning electron microscopy, 661
sensilla, 661
Tephritoidea,
agave, 430
mating behavior, 430
natural history, 430
Ulidiidae, 430
termite,
imidacloprid, 84
metabolism, 84
Reticulitermes flavipes, 84
toxicity, 84
terrestrial,
Drosophila, 247
fitness, 247
offspring number, 247
perfluorooctane sulfonic acid, 247
thermal requirements,
developmental time, 943
Planococcus ficus, 943
temperature threshold, 943
vines, 943
thermal adaptation,
acoustic behavior, 860
cicadas, 860
morphometrics, 860
systematics, 860
thinning apples,
rearing, 50
Synanthedon scitula, 50
Tomoceridae,
Appalachian Mountains, 472
Collembola, 472
phylogeny, 472
taxonomy, 472
toxicity,
imidacloprid, 84
metabolism, 84
Reticulitermes flavipes, 84
termite, 84
trade-off,
alkaloids, 371
entomopathogenic nematodes, 371
pathogen reproduction, 371
polyphagy, 371
trap-nest,
body size, 775
Megachile rotundata, 775
triose phosphate isomerase,
host strains, 283
Spodoptera frugiperda, 283
Troides aeacus,
amplified fragment length polymorphism, 1018
cytochrome oxidase I, 1018
genetic diversity, 1018
tropical forest,
cicadas, 216
diversity, 216
exuviae, 216
plantation, 216
tunnel,
deposition, 449
excavation, 449
idiothetic, 449
path integration, 449
Tuta absoluta,
coexistence, 936
Dineulophus phtorimaeae, 936
parasitoid, 936
Pseudapanteles dignus, 936
Typhlodrominae,
Ucka Nature Park, 165
Croatia, 165
faunistic survey, 165
Typhlodromus (*Typhlodromus*) *octogenipilus*, 165
Typhlodromus (*Typhlodromus*) *octogenipilus*,
Ucka Nature Park, 165
Croatia, 165
faunistic survey, 165
Typhlodrominae, 165
Typhlodromus (*Typhlodromus*) *octogenipilus*, 165

U

Ucka Nature Park,
Croatia, 165
faunistic survey, 165
Typhlodrominae, 165
Typhlodromus (*Typhlodromus*) *octogenipilus*, 165
Ulidiidae, 111, 430
agave, 430
ejaculate expulsion, 111

mating behavior, 430
natural history, 111, 430
Tephritoidea, 430
Upper Jurassic–Lower Cretaceous,
China, 866
Elateridae, 866
new genera and species, 866
Upper Jurassic–Lower Cretaceous, 866
Yixian Formation, 866
Utetes canaliculatus,
Diachasma alloeum, 908
Diachasmimorpha mellea, 908
Rhagoletis pomonella, 908
sequential speciation, 908

V

varroa,
honey bees, 146
hygiene, 146
varroa sensitive hygiene, 146
varroa mite,
Acarapis woodi, 711
Apis mellifera, 711
colony decline, 711
varroa sensitive hygiene,
honey bees, 146
hygiene, 146
varroa, 146
vector,
confidence Intervals, 827
Diaphorina, 827
pooling, 827
prevalence, 827
vegetation structure,
disturbance, 635
richness, 635
Solenopsis invicta, 635
species composition, 635
velvet ants,
biogeography, 555
desert, 555
phylogeography, 555
Sphaeropthalminae, 555
vibrational signals,
Hemiptera, 813
phytoplasma vector, 813
planthopper, 813
vines,
developmental time, 943
Planococcus ficus, 943
temperature threshold, 943
thermal requirements, 943

W

Wallacea,
mitochondrial DNA, 1025
phylogeography, 1025
wasp venom, 273, 293
cryptic, 273
cytotoxicity, 273, 293
hemocyte count, 273, 293
hemocyte viability, 273, 293
mantid, 273
natural enemy, 273
primary defenses, 273
secondary defenses, 273
whitefly,
barcoding, 196
cytochrome oxidase 1, 196
phylogenetics, 196
whole genome amplification,
Brevipalpus spp, 898
citrus leprosis, 898
SCAR markers, 898
wing margin,
butterfly, 988
Pieris rapae, 988
sensory bristle, 988
sensory scale, 988
wing morphometry,
Apis, 965
geometric morphometry, 965
honey bee, 965
Wolbachia pipientis,
agricultural pest, 1003
mitotic chromosomes, 1003
polytene chromosomes, 1003
Rhagoletis completa, 1003
worker size,
defense, 678
fire ants, 678
nest disturbance, 678
Solenopsis invicta, 678

Y

Yixian Formation,
China, 866
Elateridae, 866
new genera and species, 866
Upper Jurassic–Lower Cretaceous, 866
yolk protein,
egg development, 971
enzyme-linked immunosorbent assay, 971
Ephestia kuehniella, 971
nutrition, 971

Z

Zelus,
insect behavior, 771
parental investment, 771

α

α, β, and γ diversity
Central, Guanacaste, and Talamanca mountain ranges, 956
diversity partitioning, 956
light traps, 956